

# Ioan Raşa

Technical University of Cluj-Napoca, Romania

## Positive operators, inequalities, and stochastic convex orders

(joint work with ANA MARIA ACU)

Let  $p_{n,j}(x) := \binom{n}{j} x^j (1-x)^{n-j}$ ,  $x \in [0, 1]$ ,  $0 \leq j \leq n$ . The analytic inequality

$$\sum_{i=0}^n \sum_{j=0}^n [p_{n,i}(x)p_{n,j}(x) + p_{n,i}(y)p_{n,j}(y) - 2p_{n,i}(x)p_{n,j}(y)] f\left(\frac{i+j}{2n}\right) \geq 0,$$

valid for each convex function  $f \in C[0, 1]$ , is the simplest illustration of the results presented in this talk. It is related with the shape preserving properties of the Bernstein-Schnabl operators, see [4, Sec. 3.4]. Its first proof [6] uses stochastic convex orderings. The first *analytic* proof [1] was followed by many other proofs, in analytic or probabilistic terms, involving more general families of operators and convex functions of higher order, see [2], [5] and the references therein. The talk surveys the existing results in this area and presents some new, very recent results and problems [3].

### References

- [1] U. Abel, An inequality involving Bernstein polynomials and convex functions, *J. Approximation Theory* 222 (2017), 1-7.
- [2] U. Abel, D. Leviatan, An extension of Raşa's conjecture to  $q$ -monotone functions, *Results Math.* 75, 180 (2020).
- [3] U. Abel, D. Leviatan, I. Raşa, Relations between the Bernstein polynomials and  $q$ -monotone functions, (submitted)
- [4] F. Altomare, M. Cappelletti Montano, V. Leonessa, I. Raşa, *Markov Operators, Positive Semigroups and Approximation Processes*, Walter de Gruyter, Berlin, Munich, Boston (2014).
- [5] A. KomisarSKI, T. Rajba, Muirhead inequality for convex orders and a problem of I. Raşa on Bernstein polynomials, *J. Math. Anal. Appl.* 458(1) (2018), 821-830.
- [6] J. Mrowiec, T. Rajba, S. Wasowicz, A solution to the problem of Raşa connected with Bernstein polynomials, *J. Math. Anal. Appl.* 446(1) (2017), 864-878.