

Perturbed iterative approximation for the solution of delay differential equations

M. I. Berenguer^a, S. Lamari^b

^a Department of Applied Mathematics and Institute of Mathematics (IMAG), Escuela Técnica Superior de Ingeniería de Edificación, University of Granada, 18071 Granada, Spain

^b Department of Applied Mathematics, University of Granada, 18071 Granada, Spain
maribel@ugr.es, salmalamari@correo.ugr.es

The purpose of this work is to obtain an approximation to the solution of a delay differential equation by combining a one-step Picard-type scheme and the use of Schauder bases in a suitable Banach space. The effectiveness of the proposed method is showed with some examples.

References

- [1] A. Bellen, M. Zennaro, *Numerical methods for delay differential equations*, Numerical Mathematics and Scientific Computation, The Clarendon Press, Oxford University Press, New York (2003).
- [2] M. Dobrițoiu, M. Șerban, *Step method for a system of integral equations from biomathematics*, Applied Mathematics and Computation, 227 (2014), pp. 412–421.
- [3] D. Gámez, A. I. Garralda Guillem, M. Ruiz Galán, *Nonlinear initial-value problems and Schauder bases*, Nonlinear Analysis, 63 (2005), pp. 97–105.
- [4] B. Gelbaum, J. Gil de Lamadrid. *Bases of tensor products of Banach spaces*, Pacific Journal of Mathematics, 11 (1961), pp. 1281–1286.
- [5] A. Javid, A. Faeem, *A new iterative scheme to approximating fixed points and the solution of a delay differential equation*, Journal of Nonlinear and Convex Analysis, 21 (9) (2020), pp. 2151–2163.