Applications of fibre contraction principle to some classes of functional integral equations

Veronica Ilea[a], **Diana Otrocol**[b], Ioan A. Rus[a], Marcel A. Şerban[a]

[a] Department of Mathematics, Babeş-Bolyai University (Romania)
[b] Department of Mathematics, Technical University of Cluj-Napoca (Romania)

vdarzu@math.ubbcluj.ro, diana.otrocol@math.utcluj.ro, iarus@math.ubbcluj.ro,
mserban@math.ubbcluj.ro

By a new variant of fibre contraction principle ([4]) we give existence, uniqueness and convergence of successive approximations results for a functional integral equation. In the case of ordered Banach space, Gronwall-type and comparison-type results are also given.

References

[1] T.A. Burton, *A note on existence and uniqueness for integral equations with sum of two operators: progressive contractions*, Fixed Point Theory, 20(2019), no. 1, pp. 107–112.

[2] C. Corduneanu, *Abstract Volterra equations: a survey*, Math. and Computer Model., 32(11-13)(2000), pp. 1503–1528.

[3] V. Ilea, D. Otrocol, *Functional differential equations with maxima, via step by step contraction principle*, Carpathian J. Math., 37(2021), no. 2, pp. 195–202.

[4] A. Petruşel, I.A. Rus, M.A. Şerban, *Some variants of fibre contraction principle and applications: from existence to the convergence of successive approximations*, Fixed Point Theory, 22(2021), no. 2, pp. 795–808.